

## **RELIGION AND BODY WEIGHT IN AN UNDERSERVED POPULATION\***

**Karen H. Kim Yeary, Chan-hee Jo,  
Pippa Simpson, Jeffrey M. Gossett,  
Glenda Stein Johnson,  
Beverly J. McCabe-Sellers,  
Alma Thornton, Elaine Prewitt,  
and Bernestine B. McGee**

**Abstract:** Excess body weight is a prominent public health problem, with underserved groups bearing a disproportionate burden of disease (Ogden et al., 2006; Association AH, 2005; Hartley, 2004). Understanding body weight through examining potential predictors is an important step in addressing the current obesity epidemic. A potential predictor that has not been thoroughly examined in relationship to body weight is religion. Americans are very religious (Gallup Poll GP, 2001; Worldwide DN, 2000), with certain underserved subgroups of the population reporting greater religiosity than others. Those living in the rural South and African Americans report higher rates of church attendance and membership (Scandrett, 1996). Religion's prominence in some underserved groups that bear a disproportionate burden of the obesity epidemic may play an important role in determining body weight. Data from FOODS 2000, a representative sample of a rural, impoverished area in the lower Mississippi Delta region of the US, were analyzed (total n=1606; 787 African American and 819 Caucasian adults aged 18+). Religion was hypothesized to be associated with higher body weight; with the relationship being more pronounced in African Americans than Whites. Health behaviors (smoking, nutrition, physical activity) were expected to mediate the relationship. In Whites only, those consuming religious media at least once a week or more were 1.37 BMI units heavier than those consuming religious media less than once a week. Smoking mediated this relationship. There were no significant relationships between religion and body weight in African Americans.

**Keywords:** religion; spirituality; body weight; obesity; health disparities

---

**Karen H. Kim Yeary** is an Assistant Professor in the Department of Health Behavior and Health Education at the University of Arkansas for Medical Sciences.

Dr. Yeary conducts translational research with underserved populations to prevent chronic disease through community-based, behavioral interventions.

**Address:** Fay W. Boozman College of Public Health, University of Arkansas for Medical Sciences, 4301 West Markham #820, Little Rock, AR 72202. Ph.: (501) 526-6720, Fax: (501) 526-6709, Email: khk@uams.edu

**Chan-hee Jo,** UAMS/ACH, 1120 Marshall St, Slot 512-43, Little Rock, AR 72202. Email: JoChanhee@uams.edu

**Pippa Simpson,** Medical College of Wisconsin, Quantitative Health Sciences Translational and Biomedical Center Children's Research Institute, 8701 Watertown Plank Road Milwaukee, WI 53226-4801. Email: Psimpson@mcw.edu

**Jeffrey M. Gossett,** UAMS/ACH, 1120 Marshall St, Slot 512-43, Little Rock, AR 72202. Email: GossettJefferyM@uams.edu

**Glenda Stein Johnson,** Southern University and A&M College Ashford O. Williams Hall Post Office Box 100010. Baton Rouge, LA 70813. Email: glenda\_johnson@suagcenter.com

**Beverly J. McCabe-Sellers,** 900 S. Shackleford Rd., Suite 509, Little Rock, AR, 72211. Email: Bev.McCabe-Sellers@ars.usda.gov

**Alma Thornton,** Center for Social Research at Southern University and A&M College, Baton Rouge, LA. Email: alma\_thornton@subr.edu

**Elaine Prewitt,** Fay W. Boozman College of Public Health, University of Arkansas for Medical Sciences, 4301 West Markham #820, Little Rock, AR 72202 Email: TPrewitt@uams.edu

**Bernestine McGee,** Human Nutrition and Food Program, Southern University and A&M College, Baton Rouge, LA. Email: Bernestine\_mcghee@subr.edu

Corresponding author: Karen H. Kim Yeary.

\*Supported by: USDA, ARS project # 6251-53000-004-00D

Excess body weight is a prominent public health problem. Related to excess death (Flegal, Graubard, Williamson, & Gail, 2005) and numerous chronic diseases (Flegal, Carroll, Ogden, & Johnson, 2002), obesity and overweight are prevalent in 66% of US adults (Ogden et al., 2006). Thus efforts to understand the etiology of obesity are paramount, particularly for subgroups of the population (some minorities, rural, poor) that suffer a disproportionate burden of disease (Association AH, 2005; Hartley, 2004;

In a subsequent phone call, participants answered questions on food security (US Food Security questionnaire), household income, household participation in nutritional assistance programs, dietary habits, and religiosity information.

The sample includes 1662 respondents who completed the entire telephone survey; 1662 were weighted to represent the entire LMD. We restricted our sample to adults 18 years of age or older who reported race as either white or black. This reduced our sample size from 1662 to 1606. Due to missing data on some risk factors, the sample size is further reduced in some analyses (i.e. particularly the regression analyses).

The data collected in FOODS 2000 was weighted to adjust for nonresponse at several levels. Data were adjusted for telephone numbers with unknown eligibility, unknown residency, and for refusal to participate.

For the current analysis, the sample is representative of African American and Caucasian adults aged 18 and older of the LMD population. The sample for this study consisted of 1606 adults (Black  $n=787$ ; White  $n=819$ ).

### *Independent Variables*

Religion is a complex, multidimensional construct, and there is little consensus about how religion should be conceptualized and measured. Prior literature (Kim, Sobal, & Wethington, 2003a) was considered in deciding how religion was conceptualized in the present study. The FOODS 2000 survey contained 7 religion questions pertaining to denomination, religious attendance, religious importance, religious media, and prayer.

Denomination questions asked respondents what their religious preference was among Protestant, Catholic, Jewish, Muslim, Other, or None. A follow-up question asked about specific denomination and included choices: Baptist, Methodist, Pentecostal, Nazarene, Other, and Don't Know. Denominations were collapsed to maximize a meaningful interpretation of denomination's possible relationship with body weight. The largest religious group in the LMD area is the Southern Baptist Convention (Jones, Doty, Grammichi, et al., 2002), with approximately 60% ( $n=1662$ ) of the sample reporting a Baptist affiliation. Thus, given the predominance of those in our sample indicating a Baptist affiliation, religious denomination for this analysis was grouped into Baptist or Other.

Religious attendance is a standard component of religiosity that is often analyzed (De Vaus & McAllister, 1987). FOODS 2000 respondents were asked to choose from 7 categories (never, daily, two or more times per week, once a week, two or three times a month, once a month, less than once a month) describing how often they attend religious or spiritual services. Response categories were collapsed to produce a dichotomous variable of religious attendance: attends at least once a

week vs. attends less than once a week. Attending religious services at least once a week or more was designated as high religious attendance.

Religious importance was assessed by asking respondents how important their religious beliefs were to them using four categories: "Very Important", "Fairly Important", "Not too Important", and "Not at all Important". A dichotomous variable was created dividing those indicating that their religious beliefs were "Very Important" versus those indicating otherwise.

Religious media was a dichotomous measure constructed from averaging two single-item categorical variables. Questions included, "How often do you listen to religious radio?" and "How often do you watch religious television?" Responses ranged from 'less than once a month' to 'daily'. High religious media indicated listening or watching religious media at least once a week.

Prayer was assessed by asking respondents how often they prayed, with responses ranging from daily to less than once a month. A dichotomous variable was created, with those indicating praying daily vs. those praying less than daily.

#### *Dependent Variable*

Weight and height were self-reported in the questionnaire and used to calculate Body Mass Index (BMI). The accuracy of self-reported weight and height is generally seen as adequate for survey research (Stewart, 1982; Stunkard & Albaum, 1981). Excluding the cases with missing values and implausible outliers, the total BMI sample size available for the weighted analysis was 1582.

#### *Demographic and Control Variables*

Demographics of gender, age, race/ethnicity, education, income were assessed (Sobal, 2001). Education was categorized as less than high school education, and greater than high school education. Income was categorized into three categories: \$0-\$14,999, \$15,000-\$29,999, and \$30,000+. Age was coded as a continuous variable.

Physical functioning was assessed as a continuous variable as the physical component of the 12-item Short Form Health Survey (SF-12) (Ware, Kosinski, & Keller, 1996). An example question is, "During the past 4 weeks, have you had any of the following problems with your work or other regular daily activities as a result of your physical health: accomplished less than you would like?" Previous studies have reported Cronbach alphas of 0.72 and 0.89 (Resnick & Parker, 2001). Higher scores indicated higher physical functioning.

#### *Health Behavior Variables*

Physical activity was measured using a validated single-item question from

**Table 1:** *Summary Statistics for Variables used in the Analysis*

Variable		Prevalence (N) Or Mean/SE of Mean
Black		45% (806)
Female		54% (1019)
Household Income	\$0-\$14,999	26% (480)
	\$15,000-\$29,999	25% (410)
	\$30,000+	42% (603)
Age		Mean=45.1, SE=0.3
Education	Less than high school	24% (402)
	Greater than high school	74% (1184)
Physical Functioning		Mean=49.1, SD=0.3
Denomination	Baptist	58% (974)
	Other	42% (634)
Religious Attendance	Once a week or more	59% (970)
Religious Importance	Less than once a week	41% (635)
	Very important	81% (1328)
Prayer	Other	19% (274)
	Daily	82% (1352)
Religious Radio/TV at least once per week	Less than daily	18% (243)
	At least once a week	60% (1040)
Religious Books	Less than once a week	40% (568)
	At least once a week	60% (1016)
Smoking	Less than once a week	40% (592)
	Never	57% (936)
	Current	24% (362)
Regularly Exercised	Former	18% (297)
	Yes	70% (1067)
	No	30% (541)
Total Fat (grams)		Mean=79.0, SE=1.3
Total Calories (kcal)		Mean=2001.1, SE=25.3
Obesity		34% (556)
BMI		Mean=28.3, SE=0.2

**Table 3:** *Regression of Religion on Body Mass Index (BMI) in Caucasians<sup>1</sup>*

Religion Variable	Model 1 controlling for demographics <sup>2</sup>	Model 2 controlling for demographics + smoking	Model 3 controlling for demographics + smoking + physical activity	Model 4 controlling for demographics + smoking + physical activity + nutrition
Denomination (Baptist vs. Other)	.65 (0.50)	.87 (0.49)	.82 (0.47)	.89 (0.47)
Religious Importance (Very important vs. Other)	1.38 (0.70)	1.15 (0.67)	1.18 (0.65)	1.22 (0.66)
Religious Attendance (At least once a week or more vs. Other)	.69 (0.45)	.16 (0.41)	.30 (0.40)	.30 (0.40)
Prayer (Daily vs. Other)	-.08 (0.64)	-.40 (0.64)	-.42 (0.63)	-.44 (0.63)
Religious Media (At least once a week or more vs. Other)	1.37 (0.61)*	1.28 (0.60)*	.77 (0.75)*	1.37 (0.58)*

\*\* p&lt;.01      \* p&lt; .05

<sup>1</sup>Cells of the table represent unstandardized regression coefficients (standards errors).<sup>2</sup>Demographics controlled for included gender, age, education, and physical functioning

**DISCUSSION**

This study expands upon previous studies on religion and body weight by assessing the religion-body weight connection in an under-served population. Consistent with previous studies on religion and body weight (Ferraro, 1998; Kim, Sobal, & Wethington, 2003b), greater reported religiosity was associated with higher BMI in Whites, with every 1.0 unit increase in religious media consumption (watching religious TV or listening to religious radio) being associated with a 1.37 unit increase in BMI (~7 pounds). These relationships accounted for gender, age, education, income, and physical functioning. Also consistent with a previous study on religion and body weight (Kim, Sobal, & Wethington, 2003a), the relationship between religion and BMI in Whites was mediated in part by smoking. Physical activity and nutrition did not appear to be mediators in the relationship between religious media and BMI in Whites. There were no significant relationships between religion and body weight in African Americans.

Religion's relationship with higher body weight appears to be a consistent finding across both national and underserved samples in Whites only. To the authors' knowledge, this is the first study to specifically examine religion's relationship with body weight in African Americans. In African Americans of a rural, underserved population, religion does not appear to play a role in body weight. It is unclear why religion would not be associated with body weight in African Americans. Given the higher levels of religiosity among African Americans, there may have not been enough variance in the religion measures to assess significant relationships between religion and body weight; however the levels of religiosity were high among the Whites in our sample as well. There may have also not been enough power in our sample, although an *n* of 787 seems adequate. Historically, the black church has served as a central role in African American communities, and continues to be an influential force in the lives of many African Americans today (Eng & Hatch, 1991; Eng, Hatch, & Callan, 1985). The role of the black church in African American history may have cultivated a unique African American religiousness that is not unrelated to body weight, or not adequately captured by the study's measures.

In addition to testing religion's relationship with body weight in African Americans, this study expands upon previous studies in religion and body weight by assessing physical activity and nutrition as mediators in the relationship. In addition to the lower rates of smoking among the more religious, religion's relationship with higher body weight may be due to lower physical activity or differential dietary intake. More specifically, significant relationships between religious media and higher body weight may reflect more the physical inactivity of the religious, where religious media is a proxy for physical inactivity. However, our findings suggest that the higher body weight associated with higher religious media consumption was not due to lower physical activity. Energy and fat intake were also not mediators in the relationship, which lends evidence against religion leading to a higher energy and fat intake that contributes to weight gain. It is

unclear, then, how religious media is related to higher body weight. The activity and nutrition measures may not have been sufficiently detailed to capture the full variance of physical activity and nutrition in the sample. Given that our sample was an older one, the higher religious media and body weight may be indicative of homebound or disabled participants that were not able to regularly attend church services. However, our control of physical functioning in the analysis would have accounted for this possibility to some extent.

There are several limitations in this study. Due to the cross-sectional nature of the data, it is not possible to determine causal relationships. In addition, self-reported data may have some bias or non random measurement errors involved. Despite these limitations, however, this study expands our knowledge of the relationship between religion and body weight in a previously overlooked sample, and extends our understanding of how religion may be related to body weight. In a rural, poor, predominately minority population, religion (religious media) is related to higher body weight in Whites. Social contexts, such as religion, play an important role in how health and health behaviors are shaped. Given that religion plays a significant role in several marginalized populations that suffer a disproportionate burden of disease, better understanding this aspect of the social realm is important in understanding body weight and disparities in health.

## REFERENCES

- Association A.H. (2005). *Heart disease and stroke statistics*. Dallas, Texas: American Heart Association.
- Baecke, J.A.H., Burema, J., Frijters, J.E.R., Hautvast, J.G.A.J., & Wiel-Wetzels, W.A.M. (1983). Obesity in young Dutch adults: I, socio-demographic variables and body mass index. *International Journal of Obesity*, 7:1-12.
- Bharmal, M., & Thomas, J. (2006). 3rd. Comparing the EQ-5D and the SF-6D descriptive systems to assess their ceiling effects in the US general population. *Value in health*, 9(4):262-271.
- Briefel, R.R., Semplos, C.T., McDowell, M.A., Chien, S., & Alaimo, K. (1997). Dietary methods research in the third National Health and Nutrition Examination Survey: underreporting of energy intake. *The American journal of clinical nutrition*, 65(4 Suppl):1203S-1209S.
- Casey, P., Horton, J., Bogle, M.L., et al. (2004). Self-reported health of residents of the Mississippi delta. *Journal of Health Care for the Poor and Underserved*, 15:645-662.
- Champagne, C.M., Bogle, M.L., McGee, B.B., et al. (2004). Dietary intake in the Lower Mississippi Delta region: Results from the Foods of our Delta Study. *J Am Diet Assoc*, 104(2):199-207.
- Chatters, L.M. (2000). Religion and health: public health research and practice. *Annual Review of Public Health*, 21:335-367.
- Chen, Y., Horne, S.L., & Dosman, J.A. (1993). The influence of smoking



- cessation on body weight may be temporary. *Am J Public Health*, 83(9):1330-1332.
- Cline, K.M. & Ferraro, K.F.. Does religion increase the prevalence and incidence of obesity in adulthood? *Journal for the Scientific Study of Religion*, 45(2):269-281.
- Cohen, S., Schwartz, J.E., Bromet, E.J., & Parkinson, D.K. (1991). Mental health, stress, and poor health behaviors in two community samples. *Preventive Medicine*, 20:306-315.
- Davis, E.M., Clark, J.M., Carrese, J.A., Gary, T.L., & Cooper, L.A. (2005). Racial and socioeconomic differences in the weight-loss experiences of obese women. *American Journal of Public Health*, Sep 95(9):1539-1543.
- De Vaus, D. & McAllister, I. (1987). Gender differences in religion: A test of the structural location theory. *American Sociological Review*, 52:472-481.
- Ellison, C.G. & Levin, J.S. (1998). The religion-health connection: Evidence, theory, and future directions. *Health Education and Behavior*, 25(6):700-720.
- Eng, E., Hatch, J., Callan, A. (1985). Institutionalizing social support through the church and into the community. *Health Educ Q*, 12(1):81-92.
- Eng, E., Hatch, J.W. (1991). Networking between black churches and agencies: The lay health advisor model. *Journal of Prevention in Human Services*, 10(1):123-146.
- Ferraro, K.F. (1998). Firm believers? Religion, body weight, and well being. *Review of Religious Research*, 39:224-244.
- Flegal, K.M., Carroll, M.D., Ogden, C.L., & Johnson, C.L. (2002, Oct 9). Prevalence and trends in obesity among US adults, 1999-2000. *Journal of the American Medical Association*, 288(14):1723-1727.
- Flegal, K.M., Graubard, B.I., Williamson, D.F., & Gail, M.H. (2005, Apr 20). Excess deaths associated with underweight, overweight, and obesity. *Jama*. 293(15):1861-1867.
- Flegal, K.M., Troiano, R.P., Pamuk, E.R., Kuczmarski, R.J., & Campbell, S.M. (1995, Nov 2). The influence of smoking cessation on the prevalence of overweight in the United States. *N Engl J Med*, 333(18):1165-1170.
- Gabbe, B.J., Cameron, P.A., Graves, S.E., Williamson, O.D., & Edwards, E.R.. (2007). Preinjury status: Are orthopaedic trauma patients different than the general population? *J Orthop Trauma*, 21(4):223-228.
- Gallup Poll GP. Gallup Poll. (2001). United States: The Roper Center for Public Opinion Research.
- Gibson, R.S. (1987). Sources of error and variability in dietary assessment methods: A review. *J Can Diet Assoc*, 48:150-155.
- Goldblatt, P.B., Moore, M.E., Stunkard, A.J. (1965). Social factors in obesity. *Journal of the American Medical Association*, 192(12):97-102.
- Guenther, P.B., DeMaio, T.J., Ingwersen, L.A., & Berlin, M. (1997). The multiple-pass approach for the 24-hour recall in the Continuing Survey of Food Intakes by Individuals (CSFII) 1994-1996. *American Journal of Clinical Nutrition*, 65(Suppl):1316S.

- Hartley, D. (2004). Rural health disparities, population health, and rural culture. *American Journal of Public Health*, 94(10):1675-1678.
- Jones, D.E., Doty, S., Grammichi, C., et al. (2002). Religious congregations and membership in the United States 2000. *An enumeration by region, state and county based on data reported for 149 religious bodies*. Nashville, TN: Glenmary Research Center.
- Kim, K.H. & Sobal, J. (2004). Religion, social support, fat intake, and physical activity. *Public Health Nutrition*, 7:773-781.
- Kim, K.H., Sobal, J., & Wethington, E. (2003a). Religion and body weight. *International Journal of Obesity*. Apr 2003;27(4):469-477.
- \_\_\_\_\_. (2003b). Religion and body weight. *Int J Obes Relat Metab Disord*, 27(4):469-477.
- Koenig, H.G. (2000). Religion, well-being, and health in the elderly: The scientific evidence for an association. In: J. Thorson (Ed.), *Perspectives on spiritual well-being and aging*. Springfield, IL: Charles C. Thomas.
- Koenig, H.G., Hays, J.C., George, L.K., Blazer, D.G., Larson, D.B., & Landerman, L.R.. (1997). Modeling the cross-sectional relationships between religion, physical health, social support, and depressive symptoms. *American Journal of Geriatric Psychiatry*, 5(2):131-144.
- Koenig, H.G., Idler, E., Kasl, S., et al. (1999). Religion, spirituality, and medicine: a rebuttal to skeptics. *International Journal of Psychiatry in Medicine*, 29(2):123-131.
- Krause, N., Chatters, L.M. (2005). Exploring race differences in a multidimensional battery of prayer measures among older adults. *Sociology of Religion*, 66(1):23-43.
- Kumanyika, S. (2002a). The minority factor in the obesity epidemic. *Ethnicity and Disease*, 12(3):316-319.
- \_\_\_\_\_. (2002b). Obesity treatment in minorities. In T.A. Wadden & A.J. Stunkard (Eds.), *Handbook of obesity treatment*, pp. 416-446. New York: The Guilford Press.
- \_\_\_\_\_. (2005). Obesity, health disparities, and prevention paradigms: Hard questions and hard choices. *Preventing Chronic Disease*, 2(4):A02.
- Kumanyika, S. & Morssink, C.B. (1997). Cultural appropriateness of weight management programs. In S. Dalton (Ed.), *Overweight and weight management: The Health Professional's Guide to understanding and practice*. Gaithersburg, MD: Aspen's Publisher's Inc..
- LaPorte, R.E., Montoye, H.J., Caspersen, C.J. (1985). Assessment of physical activity in epidemiologic research: Problems and prospects. *Public Health Rep*, 100(2):131-146.
- McCullough, M.E., Hoyt, W.T., Larson, D.B., Koenig, H.G., & Thoresen, C. (2000). Religious involvement and mortality: A meta-analytic review. *Health Psychol*, 19(3):211-222.
- Mokdad, A.H., Ford, E.S., Bowman, B.A., et al. (2003, Jan 1). Prevalence of obesity, diabetes, and obesity-related health risk factors, 2001. *Journal of the American Medical Association*, 289(1):76-79.

- Navarro, A.M. (1999). Smoking status by proxy and self report: Rate of agreement in different ethnic groups. *Tob Control*, 8(2):182-185.
- Ogden, C.L., Carroll, M.D., Curtin, L.R., McDowell, M.A., Tabak, C.J., & Flegal, K.M.. (2006, Apr 5). Prevalence of overweight and obesity in the United States, 1999-2004. *Journal of the American Medical Association*, 295(13):1549-1555.
- Resnick, B., Parker, R. (2001). Simplified scoring and psychometrics of the revised 12-item Short-Form Health Survey. *Outcomes management for nursing practice*, 5(4):161-166.
- Scandrett, J.A. (1996). Health and the Black Church. *Journal of Religion and Health*, 35(3):231-244.
- Sempos, C.T., Flegal, K.M., Johnson, C.L., Loria, C.M., Woteki, C.E., & Briefel, R.R. (1993). Issues in the long-term evaluation of diet in longitudinal studies. *The Journal of nutrition*, 123(2 Suppl):406-412.
- Shatenstein, B. & Ghadirian, P. (1998). Influences on diet, health behaviours and their outcome in select ethnocultural and religious groups. *Nutrition*, 14(2):223-230.
- Sherkat, D.E. & Ellison, C.G. (1999). Recent developments and current controversies in the sociology of religion. *Annual Review of Sociology*, 25:363-394.
- Sobal, J. (2001). Social and cultural influences on obesity. In P. Bojorntorp (Ed.), *International textbook of obesity*, pp. 305-322.. New York: John Wiley and Sons.
- Stewart, A.L. (1982). The reliability and validity of self-reported weight and height. *Journal of Chronic Disease*, 35:295-309.
- Stuff, J.E., Casey, P.H., Szeto, K.L., et al. (2004). Household food insecurity is associated with adult health status. *The Journal of nutrition*, 134(9):2330-2335.
- Stuff, J.E., Horton, J.A., Bogle, M.L., et al. (2004). High prevalence of food insecurity and hunger in households in the rural Lower Mississippi Delta. *The Journal of rural health*, 20(2):173-180.
- Stunkard, A.J. & Albaum, B.A. (1981). The accuracy of self-reported weights. *American Journal of Clinical Nutrition*, 34:1593-1599.
- Taylor, C.B., Coffey, T., Berra, K., Iaffaldano, R., Casey, K., & Haskell, W.L. (1984). Seven-day activity and self-report compared to a direct measure of physical activity. *Am J Epidemiol*, 120(6):818-824.
- Taylor, R.J., Chatters, L.M., Jayakody, R., & Levin, J.S. (1996). Black and white differences in religious participation: A multisample comparison. *Journal for the Scientific Study of Religion*, 35:403-410.
- Tippett, K.S. & Cypel, Y.S. *Design and operation: The continuing Survey of Food Intakes by Individuals and the Diet and Health Knowledge Survey, 1994-1996 (Nationwide food survey report no. 96-1: accession no. PB 98-137268)*. Springfield, VA: U.S. Department of Agriculture, Agricultural Research Survey, National Information Service.

- Vartianinen, E., Seppala, T., Lillsunde, P., & Puska, P.J. (2002). Validation of self reported smoking by serum cotinine measurement in a community-based study. *Epidemiology Community Health*, 56:167-170.
- Ware, J., Jr., Kosinski, M., Keller, S.D. (1996). A 12-Item Short-Form Health Survey: Construction of scales and preliminary tests of reliability and validity. *Medical care*, 34(3):220-233.
- Wing, R.R., Hamman, R.F., Bray, G.A., et al. (2004). Achieving weight and activity goals among diabetes prevention program lifestyle participants. *Obesity Research*, 12(9):1426-1434.
- Worldwide DN. (2000). *DDB Needham Worldwide: DDB Needham Worldwide*.
- Yancey, G. (2005). A comparison of religiosity between European-Americans, African-Americans, Hispanic-Americans and Asian-Americans. *Research in the Social Scientific Study of Religion*, 16(1):83-104.